



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Genome Sequence of the Electrogenic Petroleum-Degrading *Thalassospira* sp. Strain HJ

Citation for published version:

Kiseleva, L, Garushyants, SK, Briliute, J, Simpson, DJW, Cohen, MF & Goryanin, I 2015, 'Genome Sequence of the Electrogenic Petroleum-Degrading *Thalassospira* sp. Strain HJ', *Genome announcements*, vol. 3, no. 3, e00483-15. <https://doi.org/10.1128/genomeA.00483-15>

Digital Object Identifier (DOI):

[10.1128/genomeA.00483-15](https://doi.org/10.1128/genomeA.00483-15)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Genome announcements

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Genome Sequence of the Electrogenic Petroleum-Degrading *Thalassospira* sp. Strain HJ

Larisa Kiseleva,^a Sofia K. Garushyants,^{a,b} Justina Briliute,^a David J. W. Simpson,^a Michael F. Cohen,^a Igor Goryanin^{a,c}

Biological Systems Unit, Okinawa Institute of Science and Technology, Tancha, Onna-son, Okinawa, Japan^a; A. A. Kharkevich Institute for Information Transmission Problems, Moscow, Russia^b; School of Informatics, University of Edinburgh, Edinburgh, United Kingdom^c

We present the draft genome of the petroleum-degrading *Thalassospira* sp. strain HJ, isolated from tidal marine sediment. Knowledge of this genomic information will inform studies on electrogenesis and means to degrade environmental organic contaminants, including compounds found in petroleum.

Received 8 April 2015 Accepted 14 April 2015 Published 14 May 2015

Citation Kiseleva L, Garushyants SK, Briliute J, Simpson DJW, Cohen MF, Goryanin I. 2015. Genome sequence of the electrogenic petroleum-degrading *Thalassospira* sp. strain HJ. *Genome Announc* 3(3):e00483-15. doi:10.1128/genomeA.00483-15.

Copyright © 2015 Kiseleva et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Michael F. Cohen, cohenm@sonoma.edu.

Inadvertently, through oil spills large and small, humans have amplified petroleum-degrading microorganisms on an unprecedented global scale (1). Strains isolated from shoreline sediments that regularly receive petroleum contaminants could potentially be exploited to break down petroleum in controlled environments, such as in microbial fuel cells (2).

Members of genus *Thalassospira* are Gram-negative, halophilic, and often facultative anaerobic alphaproteobacteria with the ability to utilize hydrocarbons as their sole source of carbon and energy (3). There are currently eight validly named species of *Thalassospira* (<http://www.bacterio.net>), and the genomes of seven strains are available in the GenBank database (<https://www.ncbi.nlm.nih.gov/genbank/>). Strain HJ was isolated from stagnant tidal marine sediment at Kaichu-doro Beach, Okinawa, Japan (26°19'56.1"N 127°54'0"E), in October 2013, 5 km from a large petroleum storage facility that has been in operation since the early 1970s. Strain HJ can grow on petroleum as a sole carbon source, it generates power in a microbial fuel cell and, based on *in silico* studies, and it possesses several catabolic pathways of interest (our unpublished data).

The genome of strain HJ was sequenced using the Illumina MiSeq sequencing platform (Illumina, Inc., San Diego, CA). The generated reads were trimmed and assembled *de novo* using SPAdes version 3.1.1 (4). The resulting sequence was then submitted to the NCBI Prokaryotic Genomes Automatic Annotation Pipeline for autoannotation (5). The size of the genome of strain HJ was found to be 4,269,399 bp, comprising 22 contigs, with a G+C content of 54.9%. Strain HJ contains 3,909 predicted genes, 3,838 putative coding sequences (CDS), 7 rRNAs, and 57 tRNAs.

In silico DNA-DNA hybridization (DDH) comparisons (6) between the genome of strain HJ and other accessible *Thalassospira* genome sequences in GenBank indicate the closest match to *Thalassospira profundimaris* WP0211 (7), with an estimated DDH value of 73.2% ± 3.8%, while comparisons with other *Thalassospira* strains gave values from 12.8% to 57.0% (high-scoring

segment pair length/total length, generalized linear model based). The aligned 16S rRNA gene sequences of these two organisms show 99.7% identity (1,449 bp/1,453 bp).

Nucleotide sequence accession number. The genome data have been deposited at NCBI under BioProject no. PRJNA275638 and accession no. [JYII000000000](https://ncbi.nlm.nih.gov/assembly/JYII000000000) for *Thalassospira* sp. HJ.

ACKNOWLEDGMENT

This work was supported by the Okinawa Institute of Science and Technology.

REFERENCES

- Head IM, Jones DM, Røling WF. 2006. Marine microorganisms make a meal of oil. *Nat Rev Microbiol* 4:173–182. <http://dx.doi.org/10.1038/nrmi-cro1348>.
- Wang X, Cai Z, Zhou Q, Zhang Z, Chen C. 2012. Bioelectrochemical stimulation of petroleum hydrocarbon degradation in saline soil using U-tube microbial fuel cells. *Biotechnol Bioeng* 109:426–433. <http://dx.doi.org/10.1002/bit.23351>.
- Baldani JJ, Videira SS, dos Santos Teixeira KR, Reis VM, de Oliveira ALM, Schwab S, de Souza EM, Pedraza RO, Baldani VLD, Hartmann A. 2014. The family *Rhodospirillaceae*, p 533–618. In Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryotes*. Springer Verlag, Heidelberg, Germany.
- Nurk S, Bankevich A, Antipov D, Gurevich A, Korobeynikov A, Lapidus A, Prjibelsky A, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, McLean J, Lasken R, Clingenpeel SR, Woyke T, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling genomes and mini-metagenomes from highly chimeric reads, p 158–170. In Deng M, Jiang R, Sun F, Zhang X (ed), *Research in computational molecular biology*. Springer Verlag, Heidelberg, Germany.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufio S, Li W. 2013. Prokaryotic genome annotation pipeline. The NCBI handbook [Internet], 2nd ed. National Center for Biotechnology Information, Bethesda, MD.
- Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <http://dx.doi.org/10.1186/1471-2105-14-60>.
- Lai Q, Shao Z. 2012. Genome sequence of *Thalassospira profundimaris* type strain WP0211. *J Bacteriol* 194:6957. <http://dx.doi.org/10.1128/JB.01903-12>.